

LIGHT PROPAGATION IN INHOMOGENEOUS UNIVERSE

Preliminary Results from Gravitational Lensing Experiments

PREMANA W. PREMADI

*Institute of Astronomy, Tohoku University
Sendai 981, Japan*

HUGO MARTEL

*Dept. of Astronomy, University of Texas
Austin, TX 78712, USA*

AND

RICHARD A. MATZNER

*Center for Relativity, University of Texas
Austin, TX 78712, USA*

1. Overview

Using a multi-plane lensing method that we have developed (Premadi, Martel, & Matzner 1998), we follow the evolution of light beams as they propagate through inhomogeneous universes. We use a P³M code to simulate the formation and evolution of large-scale structure. The resolution of the simulations is increased to sub-Megaparsec scales by using a Monte Carlo method to locate galaxies inside the computational volume according to the underlying particle distribution.

We consider cold dark matter models normalized to *COBE*, and perform a large parameter survey by varying the cosmological parameters Ω_0 , λ_0 , H_0 , and n (the tilt of the primordial power spectrum). This parameter survey is still in progress. Table 1 gives the values of the parameters for the models we have studied so far.

2. The Ray Shooting Experiments

For each model, we perform numerous ray-tracing experiments, propagating beams of 31^2 light rays back in time up to redshifts $z = 3$ or 5 . From the results of these experiments, we build statistics of the shear and magni-

TABLE 1. The Cosmological Parameters

Ω_0	λ_0	H_0	n	σ_8	Ω_0	λ_0	H_0	n	σ_8
1.0	0.0	65.0	0.7698	1.0	0.2	0.0	65.0	1.3188	0.5
1.0	0.0	75.0	0.7094	1.0	0.2	0.0	75.0	1.2190	0.5
1.0	0.0	65.0	0.8506	1.2	0.5	0.5	65.0	0.7808	0.8
1.0	0.0	75.0	0.7893	1.2	0.5	0.5	75.0	0.7049	0.8
0.5	0.0	65.0	0.9457	0.8	0.5	0.5	65.0	0.8807	1.0
0.5	0.0	75.0	0.8686	0.8	0.5	0.5	75.0	0.8024	1.0
0.5	0.0	65.0	1.0439	1.0	0.2	0.8	65.0	0.9326	0.6
0.5	0.0	75.0	0.9656	1.0	0.2	0.8	75.0	0.8273	0.6
0.2	0.0	65.0	1.0966	0.3	0.2	0.8	65.0	1.0702	0.8
0.2	0.0	75.0	0.9993	0.3	0.2	0.8	75.0	0.9629	0.8




Figure 1. Images of distant circular sources

fication of sources, as well as the properties of images. In Figure 1, we plot the images of distant circular sources, to illustrate interesting cases that occurred in some experiments: magnification and shear (Fig. 1a), double image (Fig. 1b), Einstein ring (Fig. 1c), and triple image (Fig. 1d).

References

Premadi, P.W., Martel, H., Matzner, R.A. (1998), *Astrophys. J.*, **Vol. 493**, pp. 10-27